

Bedarfmeldungen der Ukraine (für Spender)

(Quelle: UCPM)

| Bedarfsmeldungen der Ukraine | | |
|--|----------------------|-------|
| Güter | Bei Medizin: Details | Menge |
| Med. Equipment: | | |
| Patient monitor | | |
| Anesthesia machine | | |
| Glucometer | | |
| Electrocardiograph | | |
| Otoscope / ophthalmoscope | | |
| Defibrillator | | |
| Syringe pump | | |
| Infusion volumetric pump | | |
| Portable light source for patient examination | | |
| Vacuum aspirator | | |
| The device of artificial ventilation of lungs | | |
| Ultrasound diagnostic device | | |
| Laryngoscope with blades of different sizes | | |
| Portable breathing apparatus (Bag type "Ambu") for adults and children | | |
| Capnograph / CO2 monitor | | |
| Mobile X-ray machine | | |
| Set of surgical instruments | | |
| Specialized ambulance transport | | |

| Med. Consumables: | | |
|---|--|--|
| Stomach wash kit | | |
| Set for catheterization of the bladder | | |
| Traction tire | | |
| Portable oxygen cylinders | | |
| Central vein catheterization kits | | |
| Set for intraosseous access | | |
| Kits for lumbar puncture | | |
| Tracheostomy set | | |
| Tracheal intubation kit | | |
| Set for drainage of the pleural cavity | | |
| Immobilization tires of various types and sizes | | |
| Syringes with needles (different volumes) | | |
| Catheters for peripheral veins (different sizes) | | |
| System for transfusion of infusion solutions | | |
| Non - sterile bandage (different sizes) | | |
| Elastic fixing bandage (different sizes) | | |
| Sterile material for anti-burn measures | | |
| Sterile material for wound surface treatment | | |
| Hydrogel anti-burn bandage of different sizes | | |
| External fixation device | | |
| First Aid Equipment: | | |
| Individual sterile dressing package with elastic first aid compression component with protective moisture-resistant shell | | |
| Chemical means for a stop of bleeding (bandage hemostatic tamponing with hemostatic means) | | |
| Gel-based occlusive thoracic bandage (sticker) (with or without valve) | | |
| Thermo blanket | | |
| Mechanical means for a stop of bleeding like "CAT" | | |
| Nasopharyngeal airway (airway, tube) | | |
| Scissors for cutting clothes and shoes (atraumatic) | | |
| Film (valve) for artificial lung ventilation | | |
| Puncture needle for decompression of the pleural cavity | | |
| Hard neck collar with size adjustment | | |

| Medicine: | | |
|--|--|--|
| Aropine | injection: 1 mg (sulphate) in ampoules of 1 ml | |
| Heparin sodium | injections: 5000 IU / ml | |
| Hydroxyethylstarch | 6% infusion solution (60 mg / ml) | |
| Hydroxyethylstarch | 10% solution for infusion (100 mg / ml) | |
| Dexamethasone | injection: 4 mg / ml 1 ml in ampoules (as disodium phosphate salt) | |
| Diazepam | injections: 5 mg / ml | |
| Dopamine | injection: 40 mg / ml (hydrochloride) 5 ml in ampoules | |
| (Epinephrine /Adrenaline) | injection: 1 mg (as hydrochloride and hydrotartrate) 1 ml in ampoules, corresponding to 1.82 mg of adrenaline tartrate in 1 ml | |
| (Ethanol) | solution: 70% (denatured) | |
| (Isosorbide dinitrate) | solution for infusion: 1 mg / ml | |
| (Insulin injection (fast-acting)) | injection: 40 IU / ml in 10 ml vials | |
| (Insulin injection (fast-acting)) | injections of 100 IU / ml in 5 ml; 10 ml in a bottle | |
| Sodium lactate), complex solution of Sodium chloride + Potassium chloride + Calcium chloride + Sodium lactate) | injection of 100 U / ml and 5 ml; 10 ml in a bottle | |
| (Mannitol) | solution for infusion: 15% 100 ml, 200 ml, 400 ml in bottles, 100 ml, 250 ml, 500 ml in containers | |
| (Naloxone) | injection: 400 mcg (hydrochloride) 1 ml in ampoules | |

| | | |
|---|---|--|
| Sodium hydrogen carbonate) | injection: 1,4 % - 8,4 % | |
| Sodium chloride) | solution for injection / infusion: 0.9% isotonic (equivalent to Na + 154 mmol / l, Cl - 154 mmol / l) | |
| Complex solution of Sodium chloride + Potassium chloride + Calcium chloride | solution for infusion | |
| Neostigmine | injections: 0.5 mg / ml in 1 ml ampoules | |
| Neostigmine | injections of 2.5 mg (methyl sulfate) in 1 ml ampoules | |
| Ondansetron) | injection: 2 mg / ml 2 ml in ampoules (as hydrochloride) | |
| Ondansetron) | injection of 2 mg / ml 4 ml in ampoules (as hydrochloride) | |
| (Povidone-iodine) | solution: 10% (equivalent to 1% active iodine) | |
| (Salbutamol) | metered dose inhaler (aerosol): 100 mcg (as sulfate) per 1 dose | |
| (Furosemide) | injections: 10 mg / ml in 2 ml ampoules | |
| (Chlorhexidine) | solution: 0.05% (bigluconate) | |
| (Tranexamic acid) | injection: 50 mg / ml in an ampoule | |
| (Tranexamic acid) | 100 mg / ml in ampoules | |
| (Fresh-frozen plasma) | doses | |
| Red blood cells) | doses | |
| Cryoprecipitate | doses | |
| (Oxygen) | gas (liquid) | |
| (Oxygen) | gas | |
| (Sevoflurane) | liquid for inhalation | |
| (Ketamine) | injections: 50 mg / ml 2 ml | |
| (Ketamine) | injections: 50 mg / ml 10 ml | |
| (Ketamine) | injections: 10 mg / ml 10 ml | |

| | | |
|--------------------------------|--|--|
| (Ketamine) | injections: 10 mg / ml 50 ml | |
| (Propofol) | injections: 10 mg / ml 100 ml | |
| (Propofol) | injections: 20 mg / ml | |
| (Thiopental) | injections: 0.5 g | |
| (Thiopental) | injections: 1 g | |
| (Bupivacaine) | injections: 5 mg / ml 4 ml | |
| (Bupivacaine) | injections: 5 mg / ml 5 ml | |
| (Bupivacaine) | injections: 5 mg / ml 10 ml | |
| (Bupivacaine) | injections: 5 mg / ml 20 ml | |
| (Bupivacaine) | injections: 2.5 mg / ml in 20 ml | |
| (Bupivacaine) | injections: 2.5 mg / ml in 200 ml | |
| (Bupivacaine) | 0.5% (hydrochloride) 4 ml in ampoules | |
| (Lidocaine) | injection: 1% 2 ml in ampoules | |
| (Lidocaine) | 2% injection (hydrochloride) of 2 ml in ampoules | |
| (Lidocaine) | 5% (hydrochloride) 2 ml in ampoules | |
| (Morphine) | injection: 10 mg (sulphate or hydrochloride) 1 ml in ampoules | |
| (Fentanyl)* | solution for injection: 0.05 mg / ml in 2 ml ampoules | |
| (Acetylcysteine) | injections: 100 mg / ml 3 ml in ampoules | |
| (Acetylcysteine) | injections: 200 mg / ml in 10 ml ampoules | |
| Amoxicillin + Clavulanic acid) | tablets: 500 mg / 125 mg; | |
| Amoxicillin + Clavulanic acid) | powder for injection: 1000 mg + 200 mg | |
| (Ceftriaxone) | powder for solution for | |

| | | |
|-------------------------|---|--|
| | injection: 1 g (as sodium salt) in a vial | |
| (Cefotaxime) | powder for solution for injection: 1000 mg (as sodium salt) in a vial | |
| (Imipenem + Cilastatin) | powder for solution for injection: 500 mg (as monohydrate) + 500 mg (as sodium salt) in vials | |
| (Meropenem) | powder for injection / infusion: 1 g (as trihydrate) | |
| (Daptomycin) | powder for concentrate for solution for infusion: 500 mg | |
| (Daptomycin) | powder for solution for infusion: 500 mg | |
| (Linezolid) | solution for infusion: 2 mg / ml of 300 ml | |
| (Cefepime) | powder for solution for injection: 1000 mg; | |
| (Azithromycin) | solid oral dosage form: 500 mg | |
| (Azithromycin) | solution for oral use: 200 mg / 5 ml | |
| (Ciprofloxacin) | solution for intravenous infusion: 2 mg / ml (as hyclate) | |
| (Ciprofloxacin) | tablets: 500 mg (as hydrochloride) | |
| (Metronidazole) | injection: 500 mg per 100 ml vial | |
| (Vancomycin) | powder for solution for injection: 1000 mg (hydrochloride) in a vial | |
| (Flucytosine) | solution for infusion: 2.5 g / 250 ml | |
| (Protamine sulfate) | injections: 10 mg / ml (10,000 IU) in 5 ml | |
| Protamine sulfate) | vials | |
| Glucose) | solution for injection: 5% (isotonic) | |

| | | |
|---------------------------------|---|--|
| (Glucose) | solution for injection: 40 % | |
| (Potassium chloride) | 4% | |
| Potassium chloride) | 7.50% | |
| (Suxamethonium) | 50 mg (chloride) / ml 2 ml in ampoules | |
| Suxamethonium) | 20 mg / ml 5 ml in ampoules (suxamethonium iodide) powder for injection (chloride) in a vial | |
| (Atracurium) | 10 mg / ml (besylate) | |
| (Vecuronium) | powder for solution for injection: 10 mg (bromide) | |
| (Enoxaparin) | 20 mg / 0.2 ml; | |
| (Enoxaparin) | 40 mg / 0.4 ml; | |
| (Enoxaparin) | 100 mg / 1 ml; | |
| (Enoxaparin) | 150 mg / 1 ml | |
| (Bisoprolol) | 2.5 mg tablets | |
| (Bisoprolol) | 5 mg tablets | |
| (Bisoprolol) | 10 mg tablets | |
| (Metoprolol) | 25 mg tablets | |
| (Metoprolol) | 50 mg tablets | |
| (Metoprolol) | tablets: 100 mg | |
| Metoprolol tartrate | 1 mg / ml 5 ml | |
| Phenylephrine hydrochloride | 1% 1 ml | |
| Norepinephrine Tartrate | 2 mg / ml 4 ml | |
| Norepinephrine Tartrate | 2 mg / ml 8 ml | |
| (Furosemide) | tablets: 40 mg | |
| Amoxicillin + Clavulanic acid) | oral solution: 125 mg amoxicillin + 31.25 mg clavulanic acid / 5 ml | |
| (Amoxicillin + Clavulanic acid) | tablets: 250 mg / 62.5 mg; | |
| (Ceftriaxone) | powder for solution for injection: 250 mg (as sodium salt) in a vial | |
| (Ceftriaxone) | powder for solution for injection: 500 mg (as sodium salt) in a vial | |
| (Daptomycin) | powder for | |

| | | |
|--------------------------------|---|--|
| | concentrate for solution for infusion: 350 mg | |
| (Linezolid) | powder for oral liquid: 100 mg / 5 ml | |
| (Linezolid) | tablets: 400 mg; | |
| (Linezolid) | tablets: 600 mg | |
| (Azithromycin) | solid oral dosage form: 250 mg; | |
| (Azithromycin) | powder for oral suspension: 100 mg / 5 ml; | |
| (Azithromycin) | powder for oral suspension: 200 mg / 5 ml; | |
| (Chloramphenicol) | solid oral dosage form: 250 mg | |
| (Chloramphenicol) | solid oral dosage form: 500 mg | |
| (Chloramphenicol) | oil suspension for injection *: 0.5 g (as sodium succinate) / ml in 2 ml ampoules | |
| (Chloramphenicol) | solution for oral use: 150 mg (as palmitate) / 5 ml | |
| (Chloramphenicol) | powder for solution for injection: 0.5 g (sodium succinate) per vial | |
| (Ciprofloxacin) | solution for oral use: 250 mg / 5 ml (anhydrous) | |
| Ципрофлоксацин (Ciprofloxacin) | tablets: 250 mg (as hydrochloride) | |
| (Erythromycin) | solid oral dosage form: 200 mg; | |
| (Erythromycin) | solid oral dosage form: 250 mg; | |
| (Erythromycin) | powder for solution for oral use: | |
| (Erythromycin) | 125 mg / 5 ml (as stearate or estolate or ethyl succinate) | |
| (Metronidazole) | tablets: | |

| | | |
|--|---|--|
| Метронідазол (Metronidazole) | solution for oral use: 200 mg (as benzoate) / 5 ml | |
| (Nitrofurantoin) | solution for oral use: 25 mg / 5 ml | |
| (Clindamycin) | capsules: 150 mg (as hydrochloride) | |
| (Clindamycin) | solution for oral use: 75 mg / 5 ml (palmitate) | |
| (Vancomycin) | powder for solution for injection: 250 mg (hydrochloride) in a vial | |
| (Vancomycin) | powder for solution for injection: 500 mg (hydrochloride) in a vial | |
| (Vancomycin) | lyophilisate for solution for infusion: 500 mg per vial | |
| (Vancomycin) | lyophilisate for solution for infusion: 1000 mg per vial | |
| (Flucytosine) | capsules: 250 mg | |
| Loratadine | 10 mg | |
| (Amiodarone) | injections: 50 mg / ml 3 ml in ampoules | |
| (Verapamil) | injection: 2.5 mg (hydrochloride) / ml in 2 ml ampoules | |
| Aqua pro injectionibus/Water for injection) | solution for injection: 2 ml in ampoules | |
| Вода для ін'єкцій (Aqua pro injectionibus/Water for injection) | solution for injection: 5 ml in ampoules | |
| Aqua pro injectionibus/Water for injection) | solution for injection: 10 ml in ampoules | |
| (Activated charcoal) | solid oral dosage form: 0.25 g | |
| (Haloperidol) | injections: 5 mg / ml | |
| (Diclofenac) | injections: 25 mg / 3 ml in ampoules | |
| Декскетопрофену трометамол | 25 mg / 2 ml | |
| (Drotaverine) | injections: 20 mg / ml in 2 ml ampoules | |
| Isosorbide dinitrate) | tablets (sublingual): 5 mg | |

| | | |
|--|--|--|
| (Acetylsalicylic acid) | solid oral dosage form: 75 mg - 300 mg | |
| (Clonidine) | tablets: 0.15 mg | |
| (Clonidine) | solution for injection: 0.01% in 1 ml ampoules | |
| (Magnesium sulfate) | injections: 250 mg / ml in 5 ml | |
| (Metoprolol) | solution for injection: 1 mg / ml in ampoules | |
| (Sodium thiosulfate) | injections of 250 mg / ml of 50 ml | |
| Натрію тіосульфат (Sodium thiosulfate) | injections of 300 mg / ml in 5 ml ampoules | |
| (Sodium thiosulfate) | injections of 300 mg / ml in 10 ml ampoules | |
| (Oxybuprocaine) | eye drops, solution: 0.4% | |
| (Paracetamol) | solution for oral use: | |
| (Paracetamol) | suppositories: | |
| (Paracetamol) | solid oral dosage form: | |
| (Paracetamol) | solution for infusion: 10 mg / ml | |
| (Theophylline) | injections: 2% in ampoules | |
| Nitrous oxide) | gas | |
| (Loperamide) | solid oral dosage form: 2 mg | |
| (Methylthioninium chloride/Methylene blue) | injections: 10 mg / ml in 10 ml ampoules | |
| (Amoxicillin + Clavulanic acid) | powder for injection: 500 mg + 100 mg | |
| (Piperacillin and enzyme inhibitor) | powder for injection: 2 g (as sodium salt) + 250 mg (as sodium salt) | |
| (Piperacillin and enzyme inhibitor) | powder for injection: 4 g (as sodium salt) + 500 mg (as sodium salt) | |
| (Cefotaxime) | powder for solution for injection: 250 mg (as sodium salt) in a vial | |
| (Cefotaxime) | powder for solution for injection: 500 mg (as sodium salt) in a vial | |
| (Imipenem + Cilastatin) | powder for solution for | |

| | | |
|---------------------|--|--|
| | injection: 250 mg (as monohydrate) + 250 mg (as sodium salt) in vials | |
| (Meropenem) | powder for injection / infusion: 500 mg (as trihydrate); 1 g (as trihydrate) | |
| (Cefepime) | powder for solution for injection: 500 mg; | |
| (Cefepime) | powder for solution for injection: 2000 mg; | |
| (Chloramphenicol) | powder for solution for injection: 1 g (sodium succinate) in a vial | |
| (Gentamicin) | injections: 10 mg (as sulphate) / ml in 2 ml ampoules; | |
| (Gentamicin) | injections: 40 mg (as sulphate) / ml in 2 ml ampoules; | |
| (Erythromycin) | powder for solution for injection: 500 mg (as lactobionate) in a vial | |
| (Nitrofurantoin) | solid oral dosage form: 100 mg | |
| (Clindamycin) | injection: 150 mg (as phosphate) / ml | |
| (Glucose) | solution for injection: 10% | |
| (Glucose) | solution for injection: 50% | |
| (Calcium gluconate) | 100 mg / ml 10 ml | |
| Chlorpyramine | 20 mg / ml 1 ml | |

| Hum. Goods | | |
|---|-------------|--|
| First aid equipment | 5 K tons | |
| Batteries, power banks | 20 tons | |
| Generators | 50 K items | |
| Wheelchairs (children and adults), walkers, rollators, crutches | 100 K items | |
| Fuel (petrol, diesel, oil) | 500 K tons | |
| First aid kits | 500 K items | |
| Composition of First aid kits | | |
| Combat Application Tourniquet (CAT) | 20 000 | |
| COMBAT GAUZE Z-FOLD HEMOSTATIC | 20 000 | |
| Israeli bandage 6" | 20 000 | |
| Thermal blankets | 20 000 | |
| Decompression needles | 20 000 | |
| Trauma shears | 20 000 | |
| Tourniquets for intravenous infusion | 20 000 | |
| Venflons of various sizes: | 100 000 | |
| 14g x 0.45mm (Orange), | | |
| 16g x 0.45mm (Grey), | 100 000 | |
| 18g x 0.45mm (Green), | 300 000 | |
| 20g x 0.32mm (Pink), | 300 000 | |
| 22g x 0.25mm (Blue) | 100 000 | |
| Saline Solution. Plastic bottle. 200 ml, 250 ml, 400 ml | 100 000 | |
| Glucose 5% | 100 000 | |
| Adhesive bandage (medical plaser) | 100 000 | |
| Ringer's solution | 100 000 | |
| Voluven | 100 000 | |
| Tranexamic acid | 50 000 | |
| Analgin, diphenhydramine | 100 000 | |
| Adrenaline | 100 000 | |
| Atropine | 100 000 | |
| Dicyclic | 100 000 | |
| Vikasol (Antihemorrhagic agent) | 100 000 | |
| nabumetone | 100 000 | |
| Naubufin | 100 000 | |
| Butorphanol | 100 000 | |
| Gelofusine (Gelatin agents) | 100 000 | |
| Hydroxyethyl starch (HES/HAES) | 100 000 | |
| Backpack tactical medical MBP Coyote - Velmet | 500 | |
| Tactical medical pouches | 20 000 | |